



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2019

Usefulness of dermoscopy/dermatoscopy to improve the clinical and histopathologic diagnosis of skin cancers

Yélamos, Oriol ; Braun, Ralph P ; Liopyris, Konstantinos ; Wolner, Zachary J ; Kerl, Katrin ; Gerami, Pedram ; Marghoob, Ashfaq A

Abstract: Multiple studies have shown that dermoscopy increases the sensitivity and specificity for the detection of skin cancers compared to naked-eye examination. Dermoscopy can also lead to the detection of thinner and smaller cancers. Furthermore, dermoscopy leads to more precise selection of lesions requiring excision. In essence, dermoscopy helps clinicians differentiate benign from malignant lesions through the presence or absence of specific dermoscopic structures. Therefore, since most dermoscopic structures have direct histopathologic correlates, dermoscopy can allow the prediction of certain histologic findings present in skin cancers, thus helping select management and treatment options for select types of skin cancers. Visualizing dermoscopic structures in the ex vivo specimens can also be beneficial. It can improve the histologic diagnostic accuracy by targeted step-sectioning in areas of concern, which can be marked by the clinician before sending the specimen to the pathologist, or by the pathologist on the excised specimen in the laboratory. In addition, ex vivo dermoscopy can also be used to select tumor areas with genetic importance since some dermoscopic structures have been related to mutations with theragnostic relevance. In the second article of this continuing medical education series we review the impact of dermoscopy on the diagnostic accuracy of skin cancer, how can dermoscopy affect the histopathologic examination, and which dermoscopic features may be more relevant in terms of histological and genetic prediction.

DOI: <https://doi.org/10.1016/j.jaad.2018.07.072>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-160519>

Journal Article

Accepted Version

Originally published at:

Yélamos, Oriol; Braun, Ralph P; Liopyris, Konstantinos; Wolner, Zachary J; Kerl, Katrin; Gerami, Pedram; Marghoob, Ashfaq A (2019). Usefulness of dermoscopy/dermatoscopy to improve the clinical and histopathologic diagnosis of skin cancers. *Journal of the American Academy of Dermatology*, 80(2):365-377.

DOI: <https://doi.org/10.1016/j.jaad.2018.07.072>

Accepted Manuscript

Usefulness of dermoscopy/dermatoscopy to improve the clinical and histopathologic diagnosis of skin cancers

Oriol Yélamos, MD, Ralph P. Braun, MD, Konstantinos Liopyris, MD, Zachary J. Wolner, BA, Katrin Kerl, MD, Pedram Gerami, MD, Ashfaq A. Marghoob, MD



PII: S0190-9622(18)32728-2

DOI: [10.1016/j.jaad.2018.07.072](https://doi.org/10.1016/j.jaad.2018.07.072)

Reference: YMJD 12858

To appear in: *Journal of the American Academy of Dermatology*

Received Date: 25 February 2018

Revised Date: 1 July 2018

Accepted Date: 4 July 2018

Please cite this article as: Yélamos O, Braun RP, Liopyris K, Wolner ZJ, Kerl K, Gerami P, Marghoob AA, Usefulness of dermoscopy/dermatoscopy to improve the clinical and histopathologic diagnosis of skin cancers, *Journal of the American Academy of Dermatology* (2018), doi: <https://doi.org/10.1016/j.jaad.2018.07.072>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

-Title: Usefulness of dermoscopy/dermatoscopy to improve the clinical and histopathologic diagnosis of skin cancers

Authors: Oriol Yélamos, MD;^{1,2} Ralph P Braun, MD³ Konstantinos Liopyris, MD; ¹ Zachary J Wolner, BA; ¹ Katrin Kerl, MD; ³ Pedram Gerami, MD; ⁴ Ashfaq A Marghoob, MD; ^{1,5}

-Institution: ¹Dermatology Service, Memorial Sloan Kettering Cancer Center, New York, NY;

² Dermatology Department, Hospital Clínic, Universitat de Barcelona, Barcelona, Spain;

³Department of Dermatology, University Hospital Zürich, Switzerland;

⁴Department of Dermatology, Feinberg School of Medicine, The Robert H. Lurie Cancer Center, Northwestern University, Chicago, IL, USA

⁵Dermatology Service, Memorial Sloan Kettering Cancer Center, Hauppauge, NY

Corresponding author:

Oriol Yélamos, MD

oyelamos@gmail.com

Memorial Sloan Kettering Cancer Center

Dermatology Service

16 East 60th St. 4th Floor

New York, New York, 10022

646-888-6269

Dermatology Department

Hospital Clínic

Universitat de Barcelona

Villarroel 170, 08036 Barcelona, Spain

Funding: This research was funded in part by the NIH/NCI Cancer Center Support Grant P30

CA008748 and the Beca Excelencia Fundación Piel Sana (Dr Yélamos).

Conflicts of interest: PG has served as a consultant to Myriad Genomics, DermTech and Castle

Biosciences and has received honoraria for this. AAM has served as a consultant to 3Gen and

has received honoraria for this. The other authors do not have conflicts of interest relevant to the

current manuscript.

Word, figure, and reference count: 3485 words, 5 figures, 4 tables, 79 references

Outline:

- I. Introduction
- II. Use of dermoscopy to improve the clinical diagnostic accuracy for skin cancer detection
 - a. Advantages of dermoscopy
 - b. Limitations of dermoscopy
- III. Use of dermoscopy to assist in histological analysis
 - a. The role of the clinician
 - i. Benefits of including dermoscopic information and/or images as part of the pathology requisition form
 - ii. Advantages for the pathologist of highlighting the area/s of clinical concern within the excised lesion
 - b. The role of the pathologist
 - i. Benefits of examining the ex vivo dermoscopy before step-sectioning
- IV. Dermoscopic features with special relevance for the clinician and the pathologist
 - a. Melanocytic neoplasms
 - i. Melanoma-specific structures
 - ii. Dermoscopic structures with prognostic and therapeutic implications
- V. Dermoscopic features present in non-melanocytic neoplasms with special relevance
 - a. Dermoscopic predictors of basal cell carcinoma subtype
 - b. Dermoscopic predictors of squamous cell carcinoma subtype
- VI. Conclusion

ABSTRACT

Multiple studies have shown that dermoscopy increases the sensitivity and specificity for the detection of skin cancers compared to naked-eye examination. Dermoscopy can also lead to the detection of thinner and smaller cancers. Furthermore, dermoscopy leads to more precise selection of lesions requiring excision. In essence, dermoscopy helps clinicians differentiate benign from malignant lesions through the presence or absence of specific dermoscopic structures. Therefore, since most dermoscopic structures have direct histopathologic correlates, dermoscopy can allow the prediction of certain histologic findings present in skin cancers, thus helping select management and treatment options for select types of skin cancers. Visualizing dermoscopic structures in the ex vivo specimens can also be beneficial. It can improve the histologic diagnostic accuracy by targeted step-sectioning in areas of concern, which can be marked by the clinician before sending the specimen to the pathologist, or by the pathologist on the excised specimen in the laboratory. In addition, ex vivo dermoscopy can also be used to select tumor areas with genetic importance since some dermoscopic structures have been related to mutations with theragnostic relevance. In the second article of this continuing medical education series we review the impact of dermoscopy on the diagnostic accuracy of skin cancer, how can dermoscopy affect the histopathologic examination, and which dermoscopic features may be more relevant in terms of histological and genetic prediction.

INTRODUCTION

Dermoscopy has shown to increase the sensitivity for detecting skin cancers compared to naked-eye examination (NEE), and this increase is not occurring at the expense of a lower specificity.¹ In essence, dermoscopy leads to the biopsy of a more selective group of lesions and this is reflected in a reduction in the number of benign lesions biopsied for every skin cancer found.¹⁻³ Dermoscopy helps distinguishing benign from malignant lesions by revealing structures and patterns not visible with the NEE. Since these structures have direct histopathologic correlates (continuing medical education [CME] part 1), clinicians can more precisely predict histologic findings. In addition, visualizing dermoscopic structures in the ex vivo biopsy specimens can also be beneficial. This is particularly relevant when grossing skin cancers such as melanomas since there may be variability in tumor thickness across the tumoral area. In addition, some melanomas may be focally present within a nevus and in this scenario the correct diagnosis will be contingent upon sectioning the tissue in the appropriate plane. In fact, dermoscopy can improve grossing since only about 0.1% of a 4 mm specimen actually gets presented to the pathologist on a glass slide.⁴

Multiple methods aimed at targeting the areas to step-section have been proposed by clinicians and pathologists. Clinicians can provide descriptions or pictures of the lesion to the pathologist, or can mark the area directly on the specimen by suture, ink or punch scoring before sending it to the laboratory.⁵⁻⁹ Conversely, this marking can also be done in the laboratory by the pathologist or histo-technician since most dermoscopic structures can be identified on formalin fixed tissue,¹⁰ in a process called ex vivo dermoscopy (EVD) (figure 1).¹¹ Dermoscopy-guided sectioning may reduce the number of slides necessary to render a correct

diagnosis and this can impact cost containment in a positive manner. In addition, studies have shown that select dermoscopic structures can predict a higher degree of atypia, genetic mutations, or certain histologic subtypes when evaluating lesions suspicious for skin cancers.^{4, 12-15} Therefore, dermoscopy-guided sectioning offers an exciting opportunity for research by selecting different samples for biobanks, and allowing the detection of genetic mutations with theragnostic implications.

USE OF DERMOSCOPY TO IMPROVE THE CLINICAL DIAGNOSTIC ACCURACY FOR SKIN CANCER DETECTION

Key notes

- Dermoscopy increases the sensitivity for skin cancer detection and lowers the benign to malignant biopsy ratio.
- Limitations of dermoscopy include the learning curve and the occasional nonconformity of skin cancers to defined diagnostic criteria.
- Ultimately, the decision to biopsy a lesion requires the integration of multiple parameters including clinical information, morphologic analysis, and comparative and pattern analysis.

Advantages of dermoscopy

Since melanoma can become an aggressive tumor, maintaining a high sensitivity remains paramount and lesions clinically suspicious for malignancy should be excised. Some have suggested that it is justifiable to biopsy over 100 benign lesions in order to detect a single melanoma at an early stage.¹⁶ This aggressive approach with the aim of maintaining a high sensitivity, fails to acknowledge the harmful consequences from having such a low specificity (i.e., scarring, pain, wound infection, patient fear/anxiety) and driving up costs of health care.

Multiple meta-analyses have shown that dermoscopy improves user diagnostic accuracy for diagnosing skin cancers.^{1, 17, 18} In fact, dermoscopy enables the detection of melanomas with a sensitivity that is significantly higher than the NEE (90% vs 71%).¹ This is in part due to the misclassification of approximately 40% of melanomas as benign when using only the clinical

134 ABCDE rule.¹⁹ Dermoscopy also has a higher specificity compared to NEE (90% vs 81%).¹ This
135 results in fewer biopsies/excisions necessary to find a skin cancer (4-5:1 with dermoscopy vs 12-
136 15:1 with NEE alone^{2, 24}). One would think that reducing the benign to malignant biopsy ratio
137 (BMR) may be at the cost of detecting more advanced cancers. However, dermoscopy in fact
138 detects skin cancers at an earlier stage compared to NEE.^{3, 25-28} Thus, because dermoscopy has a
139 high sensitivity for diagnosing skin cancers, it decreases the number of unnecessary biopsies, it
140 identifies cancers earlier, and it results in a cost-effective cancer screening strategy.^{29, 30}

142 **Limitations of dermoscopy**

143 As with any diagnostic tool, dermoscopy requires training. During the training and
144 learning phase the clinicians tend to increase their sensitivity but lower their specificity. In fact,
145 during the first year after learning dermoscopy, generally the BMR increases.^{18, 34} However, after
146 gaining some experience, the clinician's specificity also increases and this eventuates into an
147 improved BMR compared to pre-dermoscopy use and NEE.³⁴

149 It is important to highlight that dermoscopy should not be used without clinical history
150 and clinical findings. The history and clinical features such as degree of firmness, and elevation
151 are important pieces of information that need to be placed in context with the dermoscopic
152 morphology. The final decision on whether to biopsy or not requires the integration of analytical
153 data (i.e., clinical ABCDE, dermoscopic features), comparative reasoning (i.e., is the lesion new
154 or changing compared to previous images), differential recognition (i.e., ugly duckling, single vs
155 multiple lesions), pattern recognition or gestalt, patient history (i.e., age, gender, comorbidities),
156 and "gut feeling".³⁵ This is important to appreciate since some skin cancers may lack specific

dermoscopic features, making them impossible to diagnose based purely on dermoscopic morphology. In addition, dermoscopic features and patterns may vary based on age, skin type, location, and extent of sun damage and this information also needs to be placed in context when interpreting dermoscopy. Despite these limitations, there is not a single study that has shown that dermoscopists perform worse as compared to non-dermoscopists.

USING DERMOSCOPY TO IMPROVE HISTOLOGICAL ANALYSIS

Key notes

- Dermoscopy can inform the pathologist on how best to process and section the tissue. This dermoscopic information can be provided to the pathologist via dermoscopic images or descriptions, and by the clinician marking the area of most concern before submitting the specimen to the laboratory.
- Since most dermoscopic features are visible after formalin fixation, dermoscopy can be used to guide step sectioning in the laboratory in a technique called ex vivo dermoscopy.
- Marking the specimen in vivo or ex vivo using dermoscopy improves the histological diagnostic accuracy and potentially reduces the costs of histologic processing.

The role of the clinician

Dermoscopy is widely used by dermatologists and non-dermatologists,^{31, 41} who are increasingly sampling a higher proportion of complex and histological equivocal lesions. For such lesions the clinical information may greatly help the pathologist in rendering the most accurate diagnosis. Clinicians can provide pathologists with clinical images, dermoscopic

images, dermoscopic descriptions, or may mark areas of interest before sending the specimen to the pathology laboratory.⁶⁻⁸ However, dermoscopy images and descriptions are only useful if sent to dermatopathologists who understand the importance of clinical-dermoscopy-histopathology correlations and who have acquired at least some knowledge about dermoscopy and their association with disease. For specimens sent to dermatopathologists who have little or no knowledge about dermoscopy, the clinician may want to consider marking the area of interest or bisecting the specimen in the plane of interest so as to insure that the area of clinical concern does in fact get sectioned. While these strategies have shown to improve the diagnostic accuracy of pathologists, currently they are rarely employed outside specialized cancer centers.⁴² As the importance of clinical-dermoscopy-pathology correlation becomes more widely appreciated, it is likely that the aforementioned methods will be more widely adopted.

Benefits of including dermoscopic information and/or images as part of the pathology requisition form:

As part of the pathology request form, an increasing number of clinicians are starting to add drawings, pictures or dermoscopic descriptors to the clinical information of lesions being submitted. This added information improves the diagnostic accuracy of histological analysis⁴³ as it directs the pathologic analysis and elicits a more careful examination of sections.⁴⁴ Ideally, pathologists should have access to clinical and dermoscopic images of complex melanocytic lesions in order to visually assess the areas of highest concern and determine orientation of step sectioning. Contrary to a common misconception, access to additional information prior to histopathologic review is not associated with diagnostic bias and actually may be helpful,⁵ even increasing the histologic interobserver agreement.⁹

Advantages for the pathologist of highlighting the area/s of clinical concern within excised lesion:

Before sending a specimen, clinicians have the opportunity to mark areas that display dermoscopic features that may have significance for the pathologist during the histopathological analysis. Clinicians can mark the specimen in multiple ways including using ink or nail varnish, placing a suture in the area of interest, scoring the area with a scalpel or a micropunch, or bisecting the specimen themselves.^{4,6-8} Irrespective of the method, the mark left by the clinician may help identify small select areas that may have been otherwise missed during the conventional step sectioning procedure (figure 2).⁴ However, clinicians must be cautioned not to damage the scored tissue to such an extent as to compromise histopathologic analysis.⁴

The role of the pathologist

Dermatopathologists routinely evaluate full-thickness skin sections in the vertical plane of a small percentage of the entire specimens volume, whereas dermoscopists evaluate the entire lesion in the horizontal plane but only to a maximum depth of the papillary dermis.⁴⁵ Therefore, both techniques are complementary and can be used in the laboratory to improve grossing. Towards this goal, pathologists need to acquire knowledge of the dermoscopy terminology and histology correlates, and dermoscopists need to acquire knowledge of histopathology (CME part 1).

Benefits of examining the ex vivo dermoscopy before step-sectioning

Dermoscopy is a diagnostic tool meant to be used in vivo directly on the patient's skin. However, dermoscopy can also be used on excised tissue because most dermoscopic features are visible even after formalin fixation.^{10, 11} This technique is called ex vivo dermoscopy (EVD),¹¹ and can help guide specimen grossing. As previously indicated (CME part 1), the colors and structures seen on dermoscopy have histopathologic correlates, which may have diagnostic and prognostic significance. For example, when grossing a pigmented melanoma, sections containing scarlike depigmentation or peppering on dermoscopy may underestimate the tumor thickness as they correspond to regression. Conversely, blue-gray areas indicate deep dermal melanocytes and thus this area will likely provide the most accurate indication of maximum tumor thickness.^{46, 47} EVD can also be useful to the pathologist when little or no clinical information is provided in the requisition form. In this scenario, reviewing the submitted specimen with a dermatoscope before grossing the lesion has shown to improve the diagnosis of ambiguous melanocytic and non-melanocytic lesions.⁴⁸⁻⁵⁰ Haspeslagh et al. found that EVD improved the detection of positive margins in keratinocyte carcinomas from ~8% to ~13%. In melanocytic lesions, this technique lead to the detection of a higher degree of atypia, ulceration, and higher mitotic rates.⁵⁰ In addition, Cabete et al. found that EVD elicited a change in the final diagnosis in ~14% of the studied cases, and EVD helped in the detection of melanomas missed with conventional step sectioning and improved the staging of melanomas.⁴⁹ In addition, EVD has shown to decrease the diagnostic turnaround time⁵⁰, a finding that indicates EVD can potentially optimize step-sectioning and reduce the costs of histopathologic processing.

EVD has some differences compared to conventional dermoscopy. While EVD clearly identifies structures containing pigment such as network, globules or streaks,^{10, 11} some structures

are less conspicuous. In EVD, there are more structureless areas and a decrease in the focus sharpness.^{10, 11, 48} Furthermore, after fixation some colors may appear enhanced (blue, brown, white), while others colors such as red may be less conspicuous due to a poor or absent visualization of blood vessels and degradation of hemoglobin after excision.^{10, 11, 51} This makes EVD challenging in amelanotic lesions. Another way to evaluate the dermoscopic features of excised specimens can be through photographs of the dermoscopic image. A dermatoscope can be attached to a camera⁵⁰ or simply attached to a smartphone (figure 1). This results in a cleaner and safer method to evaluate lesions, and provides an easy way to document the findings for future clinical, research, or academic purposes. EVD has a more important role in medium-to-large complex pigmented lesions or in wide excisions containing focal pigmented areas than on tiny specimens.¹¹ Regarding larger lesions, the same way clinicians may mark the area of concern, pathologists or histo-technicians can also mark the specimen in order to identify a given area under the microscope.⁵² In fact, it has been suggested that the marking of the specimen should be performed in the laboratory to minimize tissue loss or destruction.^{50, 52}

In the era of targeted therapies, EVD also has exciting potential in research. Several dermoscopic structures have been correlated with select genetic mutations (see below). Therefore, EVD can be used as a tool to gross specimens and select areas which can later be tested for mutations, or stored for future studies in biobanks. The dermoscopic image could then serve as an en face map of the clones present within a given lesion.⁵³

DERMOSCOPIC FEATURES WITH SPECIAL RELEVANCE FOR THE CLINICIAN AND THE PATHOLOGIST

Key points

- Several dermoscopic features are highly specific for melanoma and are called melanoma-specific structures.
- Select dermoscopic findings can predict the presence of aggressive melanomas as well as the presence of genetic mutations.
- Dermoscopy can predict the indolent vs. aggressive subtypes of keratinocyte carcinomas and may help triage lesions.
- Although relatively specific, the structures suggestive for melanocytic lesions can be encountered in non-melanocytic lesions and this may explain the discordance between the clinical/dermoscopic and the final histopathological diagnosis of many cases.

Dermoscopic features with special relevance present in melanocytic lesions

1. Melanoma-specific structures

Several dermoscopic features have been associated with a high odds ratio for melanoma when encountered in melanocytic lesions. These structures are collectively known as melanoma-specific structures. They include atypical pigment network, angulated lines, irregular streaks, negative pigment network, shiny white streaks, irregular dots and globules, irregular blotch, blue-white veil, regression structures (peppering and scarlike depigmentation), and polymorphous vessels. These features, which have well-established histopathological correlates

(CME part 1), are summarized in table I and shown in figure 3. We have described these features according to the 2016 International Dermoscopy Consensus on dermoscopic terminology.^{54, 55}

2. Dermoscopic structures associated with prognostic and therapeutic implications

A few dermoscopic structures have been shown to have clinical and prognostic significance with structures associated with melanoma arising in a nevus,^{50,62} dermal invasion,⁵⁶ Breslow >0.75mm,^{47, 57-59} mitotic activity,^{60, 61} or presence of lymph node metastases.⁶² Other dermoscopic features may also be relevant for therapeutic purposes as they are associated with genetic mutations targetable by specific therapies.^{12, 13, 15, 63} These findings are summarized in table II.

Dermoscopic predictors of basal cell carcinoma subtype

Dermoscopy can be used to predict the most common subtypes of basal cell carcinomas (BCC): nodular and superficial. Lallas et al developed an algorithm in which the most relevant discriminator between superficial and non-superficial BCC was the presence of blue-ovoid nests (in the non-superficial subtypes).⁶⁴ They also found arborizing vessels and ulceration to be associated with non-superficial BCC. Conversely, the presence of leaflike areas and serpentine vessels were associated with superficial BCC. Recently, Ahnslide et al. demonstrated that the presence of multiple small erosions in a flat lesion is predictive of superficial BCC in fair-skinned individuals.¹⁴

Thus, the findings more commonly associated with superficial BCC are serpentine vessels, multiple small erosions, flat surface and leaflike/spokewheel areas. The presence of

blue-ovoid nest, ulceration and arborizing vessels are more common in nodular BCC (figure 4, table III). This distinction can determine management since superficial BCC can potentially be treated non-surgically; streamlining the diagnostic and therapeutic process through a reduction in diagnostic biopsies. However, caution is advised since small foci of invasion cannot be excluded dermoscopically, and therefore other techniques such as reflectance confocal microscopy or optical coherence tomography can be used to identify deep tumor components.⁶⁵⁻⁶⁸ Interestingly, some dermoscopic findings have been described as therapeutic predictors. For example, the presence of ulceration in a non-treated BCC has been described to predict a good response to imiquimod, regardless of subtype.⁶⁹

Dermoscopic predictors of the squamous cell carcinoma spectrum

The lesions included in the squamous cell carcinoma (SCC) spectrum share many dermoscopic features such as scale, rosettes and vessels. However, other dermoscopic features can aid in differentiating actinic keratosis, Bowen disease, well-differentiated SCC, and invasive SCC. The findings are summarized in table IV and shown in figure 5.

CONCLUSION

In this CME series we have reviewed dermoscopic structures with their histopathology correlates and have shown that there exists an overlap between dermoscopy and histopathology. However, dermoscopy and histopathology are not equivalent. Histopathology holds an advantage over dermoscopy in that it evaluates vertical sections of tissue, which allows for the assessment of the full depth of the lesion from scanning magnification to cellular-level magnification. In addition, since histopathology is performed on paraffin-embedded tissue, it permits the use of immunohistochemical and molecular techniques to assist in diagnosis, which clearly cannot be done with dermoscopy. Conversely, unlike histology which evaluates less than 1% of the entire volume of the tumor, dermoscopy evaluates the entire surface area of the lesion in the horizontal plane, but only to the depth of the papillary dermis and not at cellular level resolution. The ability to evaluate the lesion in the horizontal plane permits the identification of certain diagnostically important structures, such as streaks, that are not commonly seen on vertical section histopathology. In addition, dermoscopy has the advantage of allowing the observer to identify colors, which may prove important in rendering a diagnosis. Other advantages of dermoscopy include the ability to evaluate the context of lesions on the skin and to monitor lesions over time to determine their biology and dynamics.

Dermoscopy is an ideal tool to enhance the diagnosis of skin cancer and in fact has a high sensitivity for the diagnosis of skin cancers while retaining a high specificity, resulting in a low BMR. In addition, dermoscopy can identify areas within a tumor that have prognostic relevance for the pathologist when performing step sectioning. Therefore, dermoscopy can improve histologic diagnostic accuracy, reduce costs of histologic processing, and offer research

355 opportunities through informed sampling of specimens for genetic testing. However, an adequate
356 knowledge of dermoscopy is required since exceptions occur. Nevertheless, dermoscopy can be a
357 bridge between clinicians and pathologists that strengthens the clinical-pathological correlation.

REFERENCES

1. Vestergaard ME, Macaskill P, Holt PE et al. Dermoscopy compared with naked eye examination for the diagnosis of primary melanoma: a meta-analysis of studies performed in a clinical setting. *Br J Dermatol* 2008;159(3):669-676.
2. Carli P, De Giorgi V, Crocetti E et al. Improvement of malignant/benign ratio in excised melanocytic lesions in the 'dermoscopy era': a retrospective study 1997-2001. *Br J Dermatol* 2004;150(4):687-692.
3. Salerni G, Teran T, Puig S et al. Meta-analysis of digital dermoscopy follow-up of melanocytic skin lesions: a study on behalf of the International Dermoscopy Society. *J Eur Acad Dermatol Venereol* 2013;27(7):805-814.
4. Merkel EA, Amin SM, Lee CY et al. The utility of dermoscopy-guided histologic sectioning for the diagnosis of melanocytic lesions: A case-control study. *J Am Acad Dermatol* 2016;74(6):1107-1113.
5. Ferrara G, Annessi G, Argenyi Z et al. Prior knowledge of the clinical picture does not introduce bias in the histopathologic diagnosis of melanocytic skin lesions. *J Cutan Pathol* 2015;42(12):953-958.
6. Firoz BF, Kennedy JS, Henning JS. Dermoscopy and suture marking as a tool to enhance diagnosis of pigmented lesions. *Dermatologic Surgery* 2008;34(8):1104-1107.
7. Braun RP, Kaya G, Masouye I et al. Histopathologic correlation in dermoscopy: a micropunch technique. *Arch Dermatol* 2003;139(3):349-351.
8. Yadav S, Vossaert KA, Kopf AW et al. Histopathologic correlates of structures seen on dermoscopy (epiluminescence microscopy). *American Journal of Dermatopathology* 1993;15(4):297-305.

9. Bauer J, Leinweber B, Metzler G et al. Correlation with digital dermoscopic images can help dermatopathologists to diagnose equivocal skin tumours. *Br J Dermatol* 2006;155(3):546-551.
10. Haspeslagh M, Vossaert K, Lanssens S et al. Comparison of Ex Vivo and In Vivo Dermoscopy in Dermatopathologic Evaluation of Skin Tumors. *JAMA Dermatol* 2016;152(3):312-317.
11. Scope A, Busam KJ, Malvey J et al. Ex vivo dermoscopy of melanocytic tumors: Time for dermatopathologists to learn dermoscopy. *Archives of Dermatology* 2007;143(12):1548-1552.
12. Bombonato C, Ribero S, Pozzobon FC et al. Association between dermoscopic and reflectance confocal microscopy features of cutaneous melanoma with BRAF mutational status. *J Eur Acad Dermatol Venereol* 2017;31(4):643-649.
13. Pozzobon FC, Puig-Butlle JA, Gonzalez-Alvarez T et al. Dermoscopic criteria associated with BRAF and NRAS mutation status in primary cutaneous melanoma. *Br J Dermatol* 2014;171(4):754-759.
14. Ahnslide I, Zalaudek I, Nilsson F et al. Preoperative prediction of histopathological outcome in basal cell carcinoma: flat surface and multiple small erosions predict superficial basal cell carcinoma in lighter skin types. *Br J Dermatol* 2016;175(4):751-761.
15. Armengot-Carbo M, Nagore E, Garcia-Casado Z et al. The association between dermoscopic features and BRAF mutational status in cutaneous melanoma: significance of the blue-white veil. *J Am Acad Dermatol* 2018;78(5):920-926.e4.
16. Aires DJ, Wick J, Shaath TS et al. Economic Costs Avoided by Diagnosing Melanoma Six Months Earlier Justify >100 Benign Biopsies. *J Drugs Dermatol* 2016;15(5):527-532.

17. Bafounta ML, Beauchet A, Aegerter P et al. Is dermoscopy (epiluminescence microscopy) useful for the diagnosis of melanoma? Results of a meta-analysis using techniques adapted to the evaluation of diagnostic tests. *Arch Dermatol* 2001;137(10):1343-1350.
18. Kittler H, Pehamberger H, Wolff K et al. Diagnostic accuracy of dermoscopy. *Lancet Oncol* 2002;3(3):159-165.
19. Salerni G, Lovatto L, Carrera C et al. Melanomas detected in a follow-up program compared with melanomas referred to a melanoma unit. *Arch Dermatol* 2011;147(5):549-555.
20. Longo C, Piana S, Marghoob A et al. Morphological features of naevoid melanoma: results of a multicentre study of the International Dermoscopy Society. *Br J Dermatol* 2015;172(4):961-967.
21. Jaimes N, Chen L, Dusza SW et al. Clinical and dermoscopic characteristics of desmoplastic melanomas. *JAMA Dermatol* 2013;149(4):413-421.
22. Menzies SW, Kreusch J, Byth K et al. Dermoscopic evaluation of amelanotic and hypomelanotic melanoma. *Arch Dermatol* 2008;144(9):1120-1127.
23. Carrera C, Segura S, Aguilera P et al. Dermoscopic Clues for Diagnosing Melanomas That Resemble Seborrheic Keratosis. *JAMA Dermatol* 2017;153(6):544-551.
24. Tromme I, Sacre L, Hammouch F et al. Availability of digital dermoscopy in daily practice dramatically reduces the number of excised melanocytic lesions: results from an observational study. *Br J Dermatol* 2012;167(4):778-786.

25. Haenssle HA, Hoffmann S, Holzkamp R et al. Melanoma thickness: the role of patients' characteristics, risk indicators and patterns of diagnosis. *J Eur Acad Dermatol Venereol* 2015;29(1):102-108.
26. Chen LL, Wei EX, Ma F et al. Rates of Dermoscopy Use for Melanoma Diagnosis in the Miami VA Medical Center. *JAMA Dermatol* 2017;153(6):602-603.
27. Argenziano G, Cerroni L, Zalaudek I et al. Accuracy in melanoma detection: a 10-year multicenter survey. *J Am Acad Dermatol* 2012;67(1):54-59.
28. Salerni G, Teran T, Alonso C et al. The role of dermoscopy and digital dermoscopy follow-up in the clinical diagnosis of melanoma: clinical and dermoscopic features of 99 consecutive primary melanomas. *Dermatol Pract Concept* 2014;4(4):39-46.
29. Tromme I, Legrand C, Devleeschauwer B et al. Cost-effectiveness analysis in melanoma detection: A transition model applied to dermoscopy. *Eur J Cancer* 2016;67:38-45.
30. Thomas L, Puig S. Dermoscopy, Digital Dermoscopy and Other Diagnostic Tools in the Early Detection of Melanoma and Follow-up of High-risk Skin Cancer Patients. *Acta Derm Venereol* 2017 Jul 5. doi: 10.2340/00015555-2719. [Epub ahead of print].
31. Argenziano G, Puig S, Zalaudek I et al. Dermoscopy improves accuracy of primary care physicians to triage lesions suggestive of skin cancer. *J Clin Oncol* 2006;24(12):1877-1882.
32. Binder M, Puespoeck-Schwarz M, Steiner A et al. Epiluminescence microscopy of small pigmented skin lesions: short-term formal training improves the diagnostic performance of dermatologists. *J Am Acad Dermatol* 1997;36(2 Pt 1):197-202.
33. Rogers T, Marino ML, Dusza SW et al. A Clinical Aid for Detecting Skin Cancer: The Triage Amalgamated Dermoscopic Algorithm (TADA). *J Am Board Fam Med* 2016;29(6):694-701.

34. Terushkin V, Warycha M, Levy M et al. Analysis of the benign to malignant ratio of lesions biopsied by a general dermatologist before and after the adoption of dermoscopy. *Arch Dermatol* 2010;146(3):343-344.
35. Marghoob AA , Scope A. The complexity of diagnosing melanoma. *J Invest Dermatol* 2009;129(1):11-13.
36. Rajadhyaksha M, Gonzalez S, Zavislan JM et al. In vivo confocal scanning laser microscopy of human skin II: advances in instrumentation and comparison with histology. *J Invest Dermatol* 1999;113(3):293-303.
37. Alarcon I, Carrera C, Palou J et al. Impact of in vivo reflectance confocal microscopy on the number needed to treat melanoma in doubtful lesions. *Br J Dermatol* 2014;170(4):802-808.
38. Gerami P, Jewell S, Morrison L et al. Fluorescence In Situ Hybridization (FISH) as an Ancillary Diagnostic Tool in the Diagnosis of Melanoma. *Am J Surg Pathol* 2009;33(8):1146-1153.
39. Bastian B, LeBoit P, Hamm H et al. Chromosomal gains and losses in primary cutaneous melanomas detected by comparative genomic hybridization. *Cancer Res* 1998;58(10):2170-2175.
40. Yélamos O , Gerami P. Predicting the outcome of melanoma: can we tell the future of a patient's melanoma? *Melanoma Manag* 2015;2(3):217-224.
41. Rosendahl C, Cameron A, McColl I et al. Dermatoscopy in routine practice - 'chaos and clues'. *Aust Fam Physician* 2012;41(7):482-487.

42. Waller JM , Zedek DC. How informative are dermatopathology requisition forms completed by dermatologists? A review of the clinical information provided for 100 consecutive melanocytic lesions. *J Am Acad Dermatol* 2010;62(2):257-261.
43. Soyer HP, Massone C, Ferrara G et al. Limitations of histopathologic analysis in the recognition of melanoma: a plea for a combined diagnostic approach of histopathologic and dermoscopic evaluation. *Arch Dermatol* 2005;141(2):209-211.
44. Bauer J, Metzler G, Rassner G et al. Dermatoscopy turns histopathologist's attention to the suspicious area in melanocytic lesions. *Arch Dermatol* 2001;137(10):1338-1340.
45. Braun R , Kerl K. Histopathologic correlation of dermoscopic structures. Available in: [https://dermoscopedia.org/Histopathologic correlation of dermoscopic structures](https://dermoscopedia.org/Histopathologic_correlation_of_dermoscopic_structures) .Last Accessed 25.8.2018
46. Malvehy J. Handbook of dermoscopy. London: Informa Healthcare; 2006.
47. Argenziano G, Fabbrocini G, Carli P et al. Clinical and dermoscopic criteria for the preoperative evaluation of cutaneous melanoma thickness. *J Am Acad Dermatol* 1999;40(1):61-68.
48. Amin K , Fraga GR. Ex vivo dermoscopy of cutaneous biopsies for melanocytic neoplasms: a retrospective review of 517 cases with histopathologic correlation. *Am J Dermatopathol* 2012;34(7):710-715.
49. Cabete J, Lencastre A , Joao A. Combined Use of Ex Vivo Dermoscopy and Histopathology for the Diagnosis of Melanocytic Tumors. *Am J Dermatopathol* 2016;38(3):189-193.
50. Haspeslagh M, Hoorens I, Degryse N et al. Pathologic Evaluation of Skin Tumors With Ex Vivo Dermoscopy With Derm Dotting. *JAMA Dermatol* 2016 Dec 28. doi: 10.1001/jamadermatol.2016.4444. [Epub ahead of print].

51. Maia M, Lellis RF, Marta AC. Ex vivo dermoscopy: Synchronic evaluation between dermatologist and dermatopathologist of melanocytic lesions. *An Bras Dermatol* 2009;84(5):553-555.
52. Haspeslagh M, Degryse N, De Wispelaere I. Routine Use of Ex Vivo Dermoscopy With "Derm Dotting" in Dermatopathology. *American Journal of Dermatopathology* 2013;35(8):867-869.
53. Malvehy J, Aguilera P, Carrera C et al. Ex vivo dermoscopy for biobank-oriented sampling of melanoma. *JAMA Dermatol* 2013;149(9):1060-1067.
54. Kittler H, Marghoob AA, Argenziano G et al. Standardization of terminology in dermoscopy/dermatoscopy: Results of the third consensus conference of the International Society of Dermoscopy. *J Am Acad Dermatol* 2016;74(6):1093-1106.
55. Kittler H, Braun R. Terminology. Available in: <https://dermoscopedia.org/Terminology> Last Accessed 25.8.2018
56. Balagula Y, Braun RP, Rabinovitz HS et al. The significance of crystalline/chrysalis structures in the diagnosis of melanocytic and nonmelanocytic lesions. *J Am Acad Dermatol* 2012;67(2):194.e191-198.
57. Argenziano G, Fabbrocini G, Carli P et al. Epiluminescence microscopy: criteria of cutaneous melanoma progression. *J Am Acad Dermatol* 1997;37(1):68-74.
58. Silva VP, Ikino JK, Sens MM et al. Dermoscopic features of thin melanomas: a comparative study of melanoma in situ and invasive melanomas smaller than or equal to 1mm. *An Bras Dermatol* 2013;88(5):712-717.
59. Stante M, De Giorgi V, Cappugi P et al. Non-invasive analysis of melanoma thickness by means of dermoscopy: a retrospective study. *Melanoma Res* 2001;11(2):147-152.

60. Ribero S, Argenziano G, Lallas A et al. Dermoscopic features predicting the presence of mitoses in thin melanoma. *J Dermatol Sci* 2017;86(2):158-161.
61. Deinlein T, Arzberger E, Zalaudek I et al. Dermoscopic characteristics of melanoma according to the criteria "ulceration" and "mitotic rate" of the AJCC 2009 staging system for melanoma. *PLoS One* 2017;12(4):e0174871.
62. Gonzalez-Alvarez T, Carrera C, Bennassar A et al. Dermoscopy structures as predictors of sentinel lymph node positivity in cutaneous melanoma. *Br J Dermatol* 2015;172(5):1269-1277.
63. Sanchez MI, Rabinovitz HS, Oliviero MC et al. Dark homogeneous streak dermoscopic pattern correlating with specific KIT mutations in melanoma. *JAMA Dermatol* 2014;150(6):633-639.
64. Lallas A, Tzellos T, Kyrgidis A et al. Accuracy of dermoscopic criteria for discriminating superficial from other subtypes of basal cell carcinoma. *J Am Acad Dermatol* 2014;70(2):303-311.
65. Cheng HM, Lo S, Scolyer R et al. Accuracy of optical coherence tomography for the diagnosis of superficial basal cell carcinoma: a prospective, consecutive, cohort study of 168 cases. *Br J Dermatol* 2016;175(6):1290-1300.
66. Iftimia N, Yélamos O, Chen C-SJ et al. Handheld optical coherence tomography-reflectance confocal microscopy probe for detection of basal cell carcinoma and delineation of margins. *J Biomed Opt* 2017;22(7):076006.
67. Kadouch DJ, Elshot YS, Zupan-Kajcovski B et al. One-stop-shop with confocal microscopy imaging versus standard care for surgical treatment of basal cell carcinoma: an

- open label, non-inferiority, randomized controlled multicenter trial. *Br J Dermatol* 2017;177(3):735-741.
68. Kadouch DJ, Leeflang MM, Elshot YS et al. Diagnostic accuracy of confocal microscopy imaging versus punch biopsy for diagnosing and subtyping basal cell carcinoma. *J Eur Acad Dermatol Venereol* 2017;31(10):1641-1648.
69. Urech M, Kyrgidis A, Argenziano G et al. Dermoscopic Ulceration is a Predictor of Basal Cell Carcinoma Response to Imiquimod: A Retrospective Study. *Acta Derm Venereol* 2017;97(1):117-119.
70. Argenziano G, Soyer HP, Chimenti S et al. Dermoscopy of pigmented skin lesions: results of a consensus meeting via the Internet. *J Am Acad Dermatol* 2003;48(5):679-693.
71. Carrera C, Marchetti MA, Dusza SW et al. Validity and Reliability of Dermoscopic Criteria Used to Differentiate Nevi From Melanoma: A Web-Based International Dermoscopy Society Study. *JAMA Dermatol* 2016;152(7):798-806.
72. Haenssle HA, Korpas B, Hansen-Hagge C et al. Seven-point checklist for dermoscopy: performance during 10 years of prospective surveillance of patients at increased melanoma risk. *Journal of the American Academy of Dermatology* 2010;62(5):785-793.
73. Tschandl P, Rosendahl C, Kittler H. Dermoscopy of flat pigmented facial lesions. *J Eur Acad Dermatol Venereol* 2015;29(1):120-127.
74. Pizzichetta MA, Talamini R, Marghoob AA et al. Negative pigment network: an additional dermoscopic feature for the diagnosis of melanoma. *J Am Acad Dermatol* 2013;68(4):552-559.
75. Salopek TG, Kopf AW, Stefanato CM et al. Differentiation of atypical moles (dysplastic nevi) from early melanomas by dermoscopy. *Dermatol Clin* 2001;19(2):337-345.

- 559 77. Zalaudek I, Giacomel J, Schmid K et al. Dermatoscopy of facial actinic keratosis,
560 intraepidermal carcinoma, and invasive squamous cell carcinoma: a progression model. J
561 Am Acad Dermatol 2012;66(4):589-597.
- 562 78. Cameron A, Rosendahl C, Tschandl P et al. Dermatoscopy of pigmented Bowen's
563 disease. J Am Acad Dermatol 2010;62(4):597-604.
- 564 79. Rosendahl C, Cameron A, Argenziano G et al. Dermoscopy of squamous cell carcinoma
565 and keratoacanthoma. Arch Dermatol 2012;148(12):1386-1392.
- 566 80. Lallas A, Pyne J, Kyrgidis A et al. The clinical and dermoscopic features of invasive
567 cutaneous squamous cell carcinoma depend on the histopathological grade of
568 differentiation. Br J Dermatol 2015;172(5):1308-1315.
- 569

570 **-ABBREVIATIONS AND ACRONYMS:**

571 AK – Actinic Keratosis

572 BCC – Basal Cell Carcinoma

573 BMR – Benign to Malignant Ratio

574 CME - Continuing Medical Education

575 EVD – Ex Vivo Dermoscopy

576 NEE – Naked-Eye Examination

577 RCM – Reflectance Confocal Microscopy










578 SCC - Squamous Cell Carcinoma

579

580 **ACKNOWLEDGEMENTS**

581 We would like to thank Dr Mary Le for her assistance in acquiring images for this CME series.

TABLES:**Table I.** Dermoscopic melanoma-specific structures and its odds ratio for melanoma








Schematic illustration	Metaphoric term	Description	Odds ratio for melanoma
	Atypical pigment network and angulated lines	Network with increased variability in the color, thickness, distribution and/or spacing of the lines. When angulated, typically they show gray color	2-9 ^{22, 70-73}
	Negative pigment network	Serpiginous interconnecting broadened hypopigmented lines that surround elongated and curvilinear globules	1.4-1.8 ^{71, 74}
	Irregular dots/globules	Clods with variability in color, size, shape, or spacing and distributed in an asymmetric fashion	1.7-4.8 ^{22, 70, 75}
	Irregular streaks (pseudopods and radial streaming)	Radial lines with bulbous projections (pseudopods) or without (radial streaming) irregularly distributed	1.5-5.8 ^{22, 70-72, 75}
	Granularity / peppering and scarlike depigmentation	Granularity: blue-gray dots Scarlike depigmentation: white area lighter than surrounding skin devoid of vessels and shiny white structures	2-18.3 ^{70-72, 75}
	Blue-whitish veil	Homogenous white blue area overlying a raised area	1.74-13 ^{70-72, 75}
	Shiny white streaks	Short white lines oriented parallel and orthogonal to each other (only seen in polarized dermoscopy)	2.5- 9.7 ^{71, 76}
	Irregular blotch	One off-centered blotch or multiple blotches	1.88-4.1 ^{22, 70-72}
	Polymorphous vessels	Simultaneous presence of multiple types of vessels	2.0-3.04 ^{22, 71, 72}

Re-printed with permission of Ralph Braun and Ashfaq Marghoob (partial content in Dermoscopedia.org).

Table II. Dermoscopic structures which have been shown to be predictors of histologic and genetic alterations when present in melanocytic lesions:








Histologic/genetic association		Dermoscopic predictors
Melanoma arising in a nevus ^{50,62}		Negative pigment network
Breslow depth >0.75 mm ^{47, 57-59}		Blue-whitish veil
		Atypical Vessels
		Abrupt cutoff
		More than 4 colors
		Streaks
		Milky red areas
		More than 2 dermoscopic structures
Dermal invasion ⁵⁶		Shiny white streaks
Mitosis in thin (<1 mm) melanoma ⁶⁰		Black color
		Peripheral streaks
Mitosis and ulceration ⁶¹		Shiny white streaks
		Blue-whitish veil
		Milky-red areas
Positive sentinel lymph node ⁶²		Blotch
		Ulceration without a pigmented network
Genetic mutations	MAPK mutations (<i>BRAF</i> , <i>NRAS</i>) ¹³	Peppering/granularity
	<i>BRAF</i> -mutated melanomas ^{12, 15}	Irregular peripheral streaks
		Ulceration
		Blue-whitish veil
	<i>BRAF</i> wild-type melanomas ¹²	Dotted vessels
	<i>KIT</i> mutations ⁶³	Dark homogeneous streaks

588 **Table III.** Dermoscopic structures associated with basal cell carcinoma subtypes:

Basal cell carcinoma subtype	Dermoscopic predictors	Schematic
Superficial basal cell carcinoma ^{14, 64}	Flat surface	
	Multiple small erosions	
	Serpentine vessels	
	Leaflike structures	
Non-superficial basal cell carcinoma ⁶⁴	Blue ovoid nest	
	Arborizing vessels	
	Ulceration	

589 Re-printed with permission of Ralph Braun and Ashfaq Marghoob (partial content in
 590 Dermoscopedia.org).
 591

Table IV. Dermoscopic structures associated with subtypes of squamous cell carcinoma subtypes.

Histologic subtype	Dermoscopic predictors	Schematic
Actinic keratosis ⁷⁷	Strawberry pattern	
Bowen disease ⁷⁸	Glomerular vessels	
	Dark dots/globules or round circles in linear arrangement	
Well-differentiated squamous cell carcinoma ^{77, 79}	White circles	
	Looped vessels	
Poorly-differentiated squamous cell carcinoma ⁸⁰	Vessels in >50% of the tumor surface	
	Diffuse arrangement of vessels	
	Bleeding	

Re-printed with permission of Ralph Braun and Ashfaq Marghoob (partial content in Dermoscopedia.org).

-FIGURE LEGENDS:

Fig 1. Evaluation of dermoscopic features on a formalin-fixed specimen (ex vivo dermoscopy) using a smartphone attached to a dermatoscope.

Fig 2. Pigmented lesion showing an overall reticular pattern with an off-centered area of atypical blue and gray dots and globules (A, upper circle) and an area with scarlike depigmentation (A, lower circle). Histopathologically, the area with atypical globules revealed a higher degree of atypia (B) as opposed to the area depicting scarlike depigmentation on dermoscopy (C). Thus, dermoscopy can be useful in identifying the areas with higher degree of atypia.

Fig 3. Dermoscopic images showing multiple melanoma-specific structures **A**, Melanoma arising in a nevus presenting with negative network (arrow) and irregular globules (arrowhead) **B**, Lentigo maligna depicting angulated lines (arrows) **C**, Invasive melanoma showing blue-whitish veil (asterisk), streaks (arrow) and irregular globules (arrowhead). **D**, Regressed melanoma presenting with atypical network (arrowheads), scarlike depigmentation and peppering (asterisk) and an irregular blotch (arrow).

Fig 4. Dermoscopic images showing dermoscopic features associated with different basal cell subtypes **A**, Superficial basal cell carcinoma presenting with an erosion (asterisk), serpentine vessels (arrowhead) and leaflike areas (arrow) **B**, Nodular basal cell carcinoma presenting with an ulcer (asterisk), arborizing vessels (arrowhead) and blue ovoid nests (arrow).

618 **Fig 5.** Dermoscopic images showing dermoscopic features associated with different squamous
619 cell subtypes **A**, Actinic keratosis presenting with an overall strawberry pattern **B**, Well-
620 differentiated squamous cell carcinoma presenting with central keratin plug (asterisk), looped
621 vessels (arrow) and white circles (arrowhead) **C**, In situ squamous cell carcinoma depicting
622 multiple linear black dots (arrowhead) **D**, Poorly-differentiated squamous cell carcinoma
623 depicting diffuse polymorphous vessels (arrowheads) and scale (asterisk).

